

What can I do about sudden cardiac arrest?



Working together to improve the diagnosis, treatment and quality of life for all those affected by arrhythmias

www.hearhythmalliance.org

Registered Charity No. 1107496

Glossary

Atria The top chambers of the heart that receive blood from the body and from the lungs. The right atrium is where the heart's natural pacemaker (sino atrial node - SAN) can be found

Automated external defibrillator (AED) An emergency lifesaving device that can be used by anyone to help restart the heart when sudden cardiac arrest occurs

Basic Life Support (BLS) A level of medical care used by members of the public to treat sudden cardiac arrest until medical help arrives

Cardiopulmonary Resuscitation (CPR) The term embraces all the procedures, from basic first aid to the most advanced medical interventions that can be used to restore the breathing and circulation in someone whose heart and breathing have stopped

Defibrillation The process in which a controlled electric shock is given to the heart, helping to re-establish a normal rhythm

Implantable Cardioverter Defibrillator (ICD) A small device implanted in a patient to allow automatic defibrillation of the heart when needed

Sudden cardiac arrest (SCA) When the heart stops beating suddenly and unexpectedly without warning

Supraventricular Tachycardia (SVT) A fast rhythm that starts in the upper chambers of the heart; less commonly associated with SCA

Ventricular Fibrillation (VF) A dangerously fast heart rhythm which causes the heart to stop pumping blood effectively. Defibrillation is needed to return the heart back to a normal rhythm. Sudden cardiac arrest can soon follow if the rhythm is not treated quickly with a shock

Ventricular Tachycardia (VT) A fast heart rhythm which can cause collapse or degenerate into VF

Wolff-Parkinson-White Syndrome (WPW) A relatively common heart condition that causes the heart to beat abnormally fast for periods of time due to an abnormal accessory electrical conduction pathway

Contents

What is an ICD/
S-ICD?

What is Sudden
Cardiac Arrest
(SCA)?

Who can be
affected by SCA?

What is the
difference between
SCA and heart
attack?

How is SCA treated?

What is an
Automated
External
Defibrillator (AED)?

The chain of
survival

Defibs Save Lives

Placing an AED in
your community

Treatments for
patients who
survive a sudden
cardiac arrest (SCA)



Important Information

This booklet is intended for use by people who wish to learn more about sudden cardiac arrest. The information within this booklet comes from research and patients' experiences. The booklet offers an explanation of sudden cardiac arrest and how it is treated. Additional information can be sourced at the websites provided.

Arrhythmia Alliance is leading a national campaign to place AEDs in local communities. For more information about Defibs Save Lives please e-mail: info@defibssavelives.org or visit www.defibssavelives.org

What is Sudden Cardiac Arrest (SCA)?

Sudden cardiac arrest (SCA) is a condition in which the heart stops beating suddenly and unexpectedly due to a malfunction in the heart's electrical system.


The cause of SCA is a life-threatening abnormal rhythm; an arrhythmia. The most common cause of SCA is ventricular fibrillation (VF).

When in VF, the heart's rhythm is so chaotic (called 'fibrillating'), that the heart trembles and is unable to effectively pump blood to the body and brain. Once the heart has entered VF, a sudden cardiac arrest may occur.

During SCA a patient will lose their pulse, stop breathing and fall unconscious. All of this can happen quickly - in fact, in a matter of seconds.

Sudden cardiac arrest strikes without warning. It does not discriminate by age, race or sex, claiming thousands of lives worldwide each year.

The heart and normal conduction



©2025 Arrhythmia Alliance

Who can be affected by SCA?

Unfortunately, anyone can suffer a sudden cardiac arrest. SCA is unpredictable and anyone regardless of their age or fitness level, can suffer SCA anywhere, at any time. Risk factors of SCA include a previous heart attack, previous SCA event, fast rhythm in the lower part of the heart, family history of SCA, structural heart abnormalities and heart failure. Although pre-existing heart disease is a common cause of cardiac arrest, many victims have never had a heart problem. Among the causes of SCA in younger people (without a previous heart attack or heart failure) are inherited or congenital arrhythmias; these include Wolf-Parkinson-White syndrome (WPW), Long QT syndrome (LQTS) and Brugada syndrome.

Wolff-Parkinson-White syndrome (WPW)

WPW results from an additional (accessory) connection pathway between the upper (atria) and lower (ventricles) chambers of the heart. This additional pathway occasionally allows very fast and unstable rhythms to develop and this can lead to SCA. These rhythm disturbances most often become apparent in teenage years or early twenties, but occasionally start earlier or later. In a patient with WPW, electrical impulses arrive at the ventricles too early disturbing the normal heart rhythm.

Occasionally this can degenerate into VF. The diagnosis is usually obvious from an electrocardiogram (ECG), although sometimes the characteristic appearances are not evident and may require additional testing to diagnose. However, many patients with WPW have few or no problems throughout their lives.

Long QT syndrome (LQTS)

Long QT is a syndrome which can cause a disturbance in the electrical system of the heart. This can predispose a person to ventricular tachycardia (VT) which can quickly degenerate into VF. The cause lies in the heart muscle cells which take slightly longer to recover from a heart beat (only by about a tenth of a second). In the presence of LQTS, SCA may be precipitated by such things as certain types of exercise, loud noises, or other sudden stimuli. Events usually occur in children or young people, but can be variable. The diagnosis is apparent from an ECG, which should also be offered to relatives of a patient shown to have Long QT syndrome (LQTS).

Brugada syndrome

Brugada syndrome is an inherited heart rhythm disorder which relates to the functioning of the heart muscle cells. It most commonly presents in people in their thirties and has a tendency to affect South East Asian populations. It can usually be diagnosed from an ECG but additional tests may be required.

Affected people suffer sudden collapse (syncope) due to VF or a very rapid form of VT called 'Torsade de Pointes'. This can lead rapidly to SCA unless treated with defibrillation.

Sudden cardiac arrest facts

- It strikes without warning; killing up to 200 people a day in the UK
- In the UK, less than 1 in 10 people survive out of hospital SCA
- It kills more people than lung cancer, breast cancer and AIDS combined
- It can happen to anyone, even young athletes
- Together with CPR, defibrillation is the only way to reestablish the heart's natural rhythm




What is the difference between SCA and heart attack?

Arrhythmias can reduce the heart's ability to work effectively, and if left untreated, a life threatening situation can arise. If the arrhythmia results in VT or VF, an extremely fast and chaotic rhythm can occur. The lower chambers of the heart quiver and the heart pumps blood ineffectively, causing SCA.

SCA: **'An electrical problem'**

SCA is different from heart attack. While heart attack is described as a 'plumbing problem', SCA is more of an 'electrical problem' that prevents the heart from functioning effectively. Heart attack can lead to SCA, but there are many other causes, such as congenital abnormalities, severe heart failure, electrocution and drug overdose.



Heart Attack: **'An plumbing problem'**

Heart Attack (the medical term is myocardial infarction or MI) occurs when part of the heart's blood supply is reduced or blocked, causing the heart muscle to become injured or die. The person is awake (conscious) and may complain of one or more of the signs and symptoms of heart attack.

Signs and Symptoms of SCA

When SCA occurs, the heart stops beating altogether. As a result, blood is no longer pumped throughout the body, including the brain. The person suddenly passes out, loses consciousness, and is lifeless. Sometimes the person may show signs of abnormal 'gaspings' which may last for several minutes.

Occasionally, SCA victims will experience 10-20 seconds of seizure-like activity (shaking of the arms and legs) at the onset of the event as the brain stops receiving blood and oxygen from the heart. The SCA victim is never awake and needs immediate help. If nothing is done, the victim will die within minutes. In fact, about 7-10% of sufferers die every minute without defibrillation.

Comotio cordis is a condition that can occur if you experience a severe blow or trauma to the chest, which can result in a SCA. Although a rare condition, it can commonly occur in young athletes participating in sporting activities, such as baseball, hockey or lacrosse. If a person is struck in the chest at a specific time in the heart rhythm cycle, the heart's electrical signal can be interrupted, resulting in the heart stopping. This can occur even when someone does not have a pre-existing heart condition. Without immediate treatment, commotio cordis can be fatal. Survival is much more likely if CPR is started quickly- approximately 40% of people survive when resuscitation begins within three minutes, compared to just 5% when it is delayed. Based on recent data, the overall survival rate for commotio cordis is estimated to be around 50-60%, likely thanks to quicker emergency response, wider availability of defibrillators, and growing public awareness of the condition.

Signs and Symptoms of Heart Attack SCA is often mistaken for a heart attack (myocardial infarction), but they differ in signs and symptoms. A heart attack may cause chest discomfort - pressure, squeezing, fullness, or pain - that can spread to the back, neck, jaw, or arms (often the left). Symptoms may last several minutes or come and go over hours or days. In contrast, SCA usually occurs without warning and leads to sudden loss of consciousness with no pulse.

Who is at higher risk of SCA? Patients with previous heart attack, heart failure or other known heart problems are at an increased risk of experiencing SCA. SCA is usually caused by VT and/or VF starting in scars or damaged areas of the heart muscle, or very occasionally due to the effects of medication that the patient may be taking. Anyone, at any age, at any time can suffer SCA.

How is SCA treated?

Cardiopulmonary resuscitation (CPR)

When someone suffers an SCA, defibrillation together with CPR is the only way to re-establish the heart's natural rhythm.


The letters CPR stand for cardiopulmonary resuscitation. The term embraces all the procedures, from basic first aid to the most advanced medical interventions that can be used to restore the breathing and circulation in someone whose heart and breathing have stopped. The 'Resuscitation Guidelines' are published by the Resuscitation Council (UK) and are available at www.resus.org.uk.

CPR alone = 9% survival

CPR + AED (automated external defibrillator) = >70% survival

What is an automated external defibrillator (AED)?

Early defibrillation is the key to surviving SCA



¹ Cummins, R.O. 1989. From concept to standard-of-care? Review of the clinical experience with automated external defibrillators. *Annals of Emergency Medicine* 18: 1269-75

- Survival rates drop 7 - 10 percent every minute without defibrillation¹
- CPR is a temporary measure that maintains blood flow and oxygen to the brain. It will not return the heart to a normal rhythm. Only defibrillation can return the heart to a normal rhythm.
- Quick action by the first person on the scene can truly make a difference in saving a life.
- Automated external defibrillators (AEDs) make early defibrillation readily available and are easy to use, even for lay people with no training.

An AED is an emergency life-saving device that can be used by anyone to help restart the heart when SCA strikes. The device is fully portable and gives the heart an electrical charge to establish a regular heartbeat. The AED will only shock when necessary.

Why is it needed?

A person who suffers sudden cardiac arrest may only be in a 'shockable rhythm' for the first few minutes; so immediate defibrillation is vital.

- CPR alone only saves 9% of people who suffer a sudden cardiac arrest.
- When used within the first three to five minutes, CPR combined with an AED can increase the chance of survival by up to 70%.

How it works

1. When turned on, the AED will instruct the user to connect the pads to the person's bare chest. All clothing above the patients' waist should be removed, including undergarments (especially underwired bras) because these can interfere with the electrical signal. The pads allow the AED to analyse their heart and determine if they require a shock.
2. If the device determines a shock is required, it will charge up in preparation to deliver a shock. The AED is completely safe as it will only deliver a charge when it determines a shockable rhythm is present.
3. When charged, the device instructs the user to ensure no one is touching the victim and then to press a button to deliver the shock. In the case of a fully automatic AED, the unit will advise the user that it will deliver the shock without further intervention.
4. When the shock is delivered, the device will instruct the user to begin CPR for a period, after which it will analyse the heart rhythm once again, advising whether a further shock to continue CPR is needed. Anyone can use an AED, no training is needed.

The Chain of Survival

Worldwide guidelines for response to sudden cardiac arrest include 'The Chain of Survival'. Quick action by the first person on-scene can truly make a difference in saving a life. The Chain of Survival represents the sequence of five events that must occur quickly to optimise a person's chance of surviving a cardiac arrest.



The five links of the chain are:

- **Early access** - Dial 999 immediately.
- **Early CPR** - Provide CPR to help maintain blood flow to the brain and organs until the arrival of the defibrillator and advanced medical care.
- **Early defibrillation** - Defibrillation is the only way to re-establish the natural rhythm following a sudden cardiac arrest.
- **Effective advanced life support** - An emergency team provides airway support, defibrillation, and intravenous medication.
- **Early advanced cardiac life support** - After initial survival of a SCA, a comprehensive management plan is made to decrease chances of further cardiac events.

Defibs Save Lives

We have worked with many community groups, clubs and individuals to help place lifesaving equipment for use in the event of an emergency.

The Defib Saves Lives campaign aims to ensure that AEDs are available in all public places. They must be accessible to everyone whatever the time of day. AEDs should ideally be housed in secure, weatherproof, heated cabinets on the exterior walls of buildings such as post offices, village halls, sports centres and other visible places in the community.

If you are a school or a sports team, you may decide to purchase two AEDs; one to be housed in an external cabinet at your club and a portable AED to take on trips away. A public-access AED in an external, heated and secure cabinet is the most versatile installation to ensure that an AED is always available when needed. These are ideal for high streets, shopping centres, sports stadiums, exhibition centres and concert venues.

If an AED is designed for portable use, it might still benefit from being stored in a visible, accessible location, such as a dedicated indoor cabinet. This prevents AEDs from being hidden away or forgotten in desk draws or locked offices.



Placing an AED in your community

Along with advice and guidance on purchasing your defibrillator, we offer a great range of fundraising materials and information. These include training resources and raising awareness in your community about SCA and the importance of knowing where and how to use your nearest AED. We tailor each package to suit your needs and community. We also provide guidance and advice on whether to choose a locked external cabinet, hanging hooks for the AED if internal, and rucksacks for carrying your AED and first aid kit (Grab and Go - ideal for sports and outdoor activities). All of which can be ordered through us.

Upon receipt of the defibrillator and educational package you will also receive instructions for registering your AED on the national database 'The Circuit'.



We offer a variety of AED packages to suit different communities.

For more information, please contact info@defibssaveslives.org or call our patient helpline on 01789 867501.

Treatments for patients who survive a sudden cardiac arrest (SCA)


Patients who survive a SCA or who are diagnosed as being at risk of SCA can be treated in a number of ways. Many will be implanted with an implantable cardioverter defibrillator (ICD), a device like a pacemaker which is placed beneath the skin (usually on the upper chest wall) and has wires connecting it to the heart. This device continually monitors the heart and will deliver a shock to defibrillate the heart if needed. Being fully implanted and completely automatic, the patient is able to lead a normal life with few limitations, safe in the knowledge that the ICD will respond immediately if required.

Some patients may only need to take medication alone, or in addition to an ICD. Occasionally some causes of SCA (such as WPW) can be treated by a curative procedure whereby the additional electrical pathway within the heart is destroyed by a small burn inside the heart, using a technique known as catheter ablation (see the Arrhythmia Alliance catheter ablation for cardiac arrhythmias booklet).



All patients who have survived a SCA should be reviewed by a cardiac electrophysiologist (a doctor specialising in heart rhythm disturbances) in order to determine how best to prevent further events and to consider whether family members need to be screened.

Following recent studies, it is now known that cardiac arrest survivors who are more mindful report fewer symptoms of depression, anxiety or post-traumatic stress disorder following the event. Mindfulness-based interventions such as meditation or cognitive behavioural therapy can help in decreasing these symptoms and have a positive impact on cardiovascular disease and risk.



Working together to improve the diagnosis, treatment and quality of life for all those affected by arrhythmias



+44 (0)1789 867 501

info@defibssavelives.org

www.hearhythmalliance.org

Registered Charity No. 1107496

©Arrhythmia Alliance
©AF Association
©STARS

Published 2005
Reviewed March 2025



“The information in the booklet gave me some reassurance following my husband’s SCA. I am so thankful that the defibrillator was on hand at the leisure centre”

Alison, Manchester

To view our patient resources, scan the QR code below:



Please remember that this publication provides general guidelines only. Individuals should always discuss their condition with a healthcare professional. If you would like further information or would like to provide feedback, please contact Arrhythmia Alliance.

Acknowledgments: Arrhythmia Alliance would like to thank all those who helped in the development and review of this publication. Particular thanks are given to Dr Charlotte D’Souza

Founder and Trustee:
Trudie Lobban MBE, FRCP (Edin)