

Warfarin and diet

This factsheet is intended to help those affected by atrial fibrillation understand more about the interaction between warfarin and certain types of food, to help patients maintain a stable INR level.

Many people with atrial fibrillation may find that warfarin (also known as Coumadin or Jantoven in the US) therapy is recommended to thin the blood and reduce their risk of stroke.

Warfarin is a very effective medication. However, it does require regular monitoring to assess how thin it makes the blood. It works by interfering with how the liver uses the vitamin K taken in our diet.

Warfarin's effectiveness, is thus affected, by the amount of vitamin K in the diet. If the diet is reasonably consistent, then the amount of vitamin K in your diet will be matched by the warfarin dose.

If your diet contains foods rich in vitamin K, there is no need to change your diet or restrict these foods from normal levels of consumption.

If you wish to make a more sustained change to your diet, such as eating green leafy vegetables due to their rich calcium content, you should discuss this with your anticoagulation clinic or your doctor, as they may wish to monitor your blood a little more frequently to ensure the warfarin continues to work effectively.

If you change your diet to contain foods that are richer in vitamin K then you will find you require more warfarin. If the diet changes to contain more foods that are low in vitamin K then your warfarin dose may need to be reduced.

A simple rule is that leafy green vegetables tend to be rich in vitamin K and root vegetables, fruits and cereals tend to be low in vitamin K.

It is always best to discuss your dietary needs with a qualified nutritionist or medical professional.

Foods thought to affect warfarin control:

- Asparagus
- Avocado
- Green beans
- Blackberries
- Blueberries
- Broccoli
- Brussels sprouts
- Cabbage
- Chicory
- Collard greens
- Kale
- Kiwi fruit
- Lettuce
- Mungo beans
- Mustard greens
- Peas
- Pine nuts
- Raisins
- Sugar snap peas
- Soybeans
- Spinach
- Swiss chard
- Watercress

It is important to keep a healthy and relatively consistent diet. You should not try to change your International Normalized Ratio (INR) level by regularly altering your diet as this may lead to poor INR control.

Within our normal diet other factors that can interact with warfarin include:

Alcohol: This affects the way that the body, particularly the liver, functions. Taken in a sensible manner and in moderation it is not totally contraindicated. However, periods of marked excess must be avoided.

Natural health food products: These are popular additions to the diet and may potentially affect warfarin therapy, or they may interact with the natural clotting of the blood.

The following is not an extensive list. However, any addition of this kind of product to the diet should be discussed with your doctor or anticoagulation clinic:

- American ginseng
- Asian ginseng
- Chinese angelica
- Chinese ginseng
- Chinese wolfberry
- Chondroitin plus glucosamine
- Coenzyme o10
- Devil's claw
- Dong quai
- Essence of tortoise shell
- Feverfew
- Fenugreek together with boldo
- Fish oil supplements that contain eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)

- Ginkgo biloba
- Ginseng
- Green tea
- Horse chestnut
- Japanese ginseng
- Korean ginseng
- Lycium barbarum
- Methyl salicylate (used on the skin)
- Papaya extract
- St. John's Wort
- Vitamin A
- Vitamin K
- Wintergreen (used on the skin)

Although these lists look restrictive and extensive, the key is to eat a sensible diet. Take the warfarin at approximately the same time each day. If you wish to make major changes to your diet or start taking a 'health supplement', discuss this with your doctor or the clinic which monitors your warfarin therapy. In general, they will not restrict your choices but may wish to increase the level of monitoring for a period as you make changes. Once your blood levels are seen to be stable again, then you will return to your normal monitoring intervals.

If you become ill it is important to remember that this may affect your INR level, and it would be advisable for you to have more frequent blood tests to be certain that your INR level is stable.

These lists are not intended to be comprehensive and we are aware that different authorities have different opinions on how certain foods may or may not affect warfarin. This sheet is to act as a simple guide to patients, carers or health care professionals to assist in this complex area.

Acknowledgements: AF Association would like to thank all those who helped in the development of this publication. In particular, thanks is given to Dr Charlotte D'Souza and Francesca Lobban.