

## AFib-related stroke

This information sheet is intended for people with atrial fibrillation (also referred to as AFib or AF) and their carers, and gives a brief outline of strokes and AFib-related strokes, their symptoms, and what to do in an emergency.

### AFib and stroke

AFib is a type of irregular heart rhythm. It means that your heart may not be pumping as well as it should. As a result, blood clots are more likely to form in your heart, increasing your risk of having a stroke.

A person having no other problems apart from suffering from AFib over the age of 65, has a fivefold increase in their risk of suffering from an AFib-related stroke compared with someone of the same age without AFib. This risk of stroke is present even if the AFib comes and goes (so called Paroxysmal Atrial Fibrillation or pAF). The risk is there even if the AFib does not cause any symptoms or has not been detected yet.

**Patients who suffer from AFib but take no anticoagulation medication have a 35% chance of eventually suffering from an AFib-related stroke.**

Although appropriate anticoagulation can reduce this risk, there remains a small risk of suffering from an AFib-related stroke. It is important to recognize the signs and what you, or someone with you, should do if a stroke occurs.

### What is a stroke?

Stroke is where the oxygen supply to an area of the brain has been disrupted. The two main kinds of stroke are:

**Ischemic stroke:** This is the most common form of stroke, and it occurs when the blood supply to the brain is blocked by a clot or a narrowing of the artery. The brain becomes starved of oxygen, so the brain cells suffocate. Anticoagulants are prescribed to safeguard patients with AFib against this kind of stroke.

**Hemorrhagic stroke:** This happens when one of the blood vessels in the brain leaks and the raised blood pressure forces blood into the brain tissue (an 'intracranial bleed'). This causes cells to die because of direct trauma of the blood flow.

### AFib-related stroke risk

The following scoring system ('CHA<sub>2</sub>DS<sub>2</sub>-VA') can be used to calculate an individual's risk of an AFib-related stroke. In this system, a score of 1 or more suggests that an anticoagulant should be considered or recommended, to reduce their risk of an AFib-related stroke.

CHA <sub>2</sub> DS <sub>2</sub> -VA scoring system	
Risk factor	Points
Chronic Heart Failure	1
Hypertension	1
Age 65-74 years	1
Age ≥ 75 years	2
Diabetes	1
Stroke or previous TIA	2
Vascular heart disease	1
CHA <sub>2</sub> DS <sub>2</sub> -VA Score	

### Preventing an AFib-related Stroke

If you have AFib, you should discuss your AFib-related stroke risk with your doctor and ensure preventative measures are in place to lower your risk of this happening. Maintaining a healthy lifestyle by eating a balanced diet, exercising regularly, not smoking, and minimizing alcohol intake to improve your overall health and reduce AFib-related stroke risk.

Medical treatment options include anticoagulation which increase the time it takes for blood to clot, thus reducing the risk of an AFib-related stroke. Anticoagulants include warfarin and non-vitamin K antagonist oral

endorsed by:



Founder & CEO: Trudie Lobban MBE, FRCP (Edin)  
Executive Director: Francesca Calahan

© AF Association

Published January 2010 | Reviewed December 2025



anticoagulants such as apixaban, dabigatran, edoxaban and rivaroxaban. There are risks associated with anticoagulants such as excessive bleeding if you fall or hurt yourself, but in most people the benefits outweigh the risks.

### Symptoms or signs of a stroke - The FAST test

The symptoms of a stroke can be recognized by using the FAST test, which is the test also used by paramedics to diagnose a stroke.

FAST is a useful and simple tool you can use to diagnose a stroke:

**Facial weakness**

- Can the person smile?
- Has their mouth or eye drooped?

**Arm weakness**

- Can the person raise both arms?

**Speech problems**

- Can the person speak clearly and understand what you say?

**Time to call 911**

- If a person fails any of the tests above, it is time to call for help by calling 911.

Stroke is an **emergency**. Receiving hospital assistance quickly can limit further damage and can help someone make a full recovery. **If not treated appropriately, a stroke can result in permanent disability and even be fatal.**

### If the symptoms settle quickly

In a situation where the symptoms of the FAST test seem to disappear quickly and normal body function is returned, this may suggest that a 'mini-stroke' (a 'Transient Ischemic Attack' or TIA) has occurred. Although immediate help may not be required, the sufferer should attend their GP or local hospital within 24 hours to be referred to a specialist stroke service. These mini-strokes can be signs

that there is a risk of an imminent major stroke and urgent medication changes may be required.

### Outcome

The cause of the stroke will determine the treatment options. Recovery may be complete after a short period of time, but long term recovery and rehabilitation may be necessary depending on the severity of the stroke.

If you have AFib, an anticoagulant should be considered after an ischemic stroke at any age. If you were already taking an anticoagulant at the time of your stroke, you may require additional preventative treatment such as a left atrial appendage occlusion device and should be referred to a specialist for assessment.

**STROKE IS A MEDICAL EMERGENCY  
AND ASSISTANCE SHOULD BE  
SOUGHT WITHOUT DELAY!**

### Further information

The American Stroke Association is a useful source of information, advice and support for those who have suffered a stroke or are caring for someone affected by a stroke.

**Website:** [www.stroke.org](http://www.stroke.org)

Please also see the AF Association Preventing AFib-related stroke: anticoagulation booklet.

To view our patient resources, scan the QR code below:



**Acknowledgments:** AF Association would like to thank all those who helped in the development and review of this publication. Particular thanks are given to Dr Matthew Fay, Prof Gregory Lip, Dr Steve Murray, Mr Sotiris Antoniou, Dr Charlotte D'Souza, Dr Nikhil Ahluwalia, Dr Benjamin Steinberg and Francesca Calahan.

*endorsed by:*



**Founder & CEO:** Trudie Lobban MBE, FRCP (Edin)  
**Executive Director:** Francesca Calahan

© AF Association

Published January 2010 | Reviewed December 2025