

Inherited Arrhythmias and Genetic Testing



Working together to improve the diagnosis, treatment and quality of life for all those affected by arrhythmias

www.heartrhythmalliance.org

Glossary

Inherited Cardiac Conditions (ICCs) Conditions that can be passed down in families via your genes

Electrocardiogram (EKG) is a simple test that records the heart's rhythm and rate

Long QT Syndrome (LQTS) An inherited condition where there are problems with the electrical activity of the heart

Brugada Syndrome A rare inherited heart rhythm disorder, in which the patient is at a risk of developing a fast heart rhythm

Catecholaminergic Polymorphic Ventricular Tachycardia (CPVT) A rare inherited arrhythmia caused by an abnormality of ion channels, leading to abnormal heartbeats

Important Information

This information leaflet has been produced for people who have been diagnosed with an Inherited Cardiac Condition and their families. Its aim is to outline the process and implications of genetic testing and to give you an idea of what to expect from your genetic counselling appointment.

Contents

What is an inherited cardiac condition (ICC)?

What is genetic testing?

Genetic screening

Pre-symptomatic genetic testing

What is genetic counselling?

How is the genetic test done?

Where can I get more information?

Long QT Syndrome

Brugada Syndrome

CPVT

What is an inherited cardiac condition (ICC)?

Inherited Cardiac Conditions (ICCs), for example Long QT Syndrome (LQTS) and Brugada Syndrome, are conditions that can be passed down in families via your genes. Genes act as instructions telling our bodies how to grow, develop and function. There are many genes known to be involved with ICCs. A change, or 'spelling mistake', in one of these genes can cause someone to be affected by, or be at risk of, an ICC.

The way these 'spelling mistakes' are inherited can vary, but there is most commonly a 50:50 Co or one in two chance that a close blood relative (child, parent, sister or brother) also has the 'spelling mistake' themselves, and therefore has the risk of developing the ICC too. Most people with ICCs do not suffer troublesome ongoing symptoms. However a small number of people are at risk of dangerous heart problems and it is important to have regular heart checks, as many of these problems are treatable or preventable.

The heart and normal conduction



© 2012 Arrhythmia Alliance

What is genetic testing?

A genetic test is a type of medical test that is used to try and identify the gene change, or 'spelling mistake' that can cause an ICC. For example:

A correctly 'spelled gene: A gene with a 'spelling mistake':

THE CAT SAT MAT THE CAT TSA TMA T

Within the genes that are associated with causing ICCs, there are a large number of different gene changes that can cause the condition. It is not always possible, using current technology, to find the 'spelling mistake'. If this is the case, your sample may be kept and stored as it may be possible to do additional testing in the future. Even if the gene change causing the condition in your family is found, your sample may still be stored for further testing if and when medical knowledge increases and new technologies become available.

You may be asked for your written consent before you give a sample, and your permission will also be gained before any results are shared for the purposes of helping other family members who are also considering genetic testing.

Are there different types of genetic tests?

There are two different types of genetic tests: 'genetic screening' and 'presymptomatic' or 'familial' testing. The first affected person in the family to come forward for testing will be offered a genetic screen; if a causative spelling mistake is found, then their relatives may be offered a familial test. The process involved and implications to consider are quite different for each type of test.

Genetic screening

Genetic screening can be offered to someone who has been diagnosed with an ICC by their specialist heart doctor (cardiologist) based on the results of clinical tests (like EKG and echocardiography) and examination.

The aim of genetic screening is to try and identify the genetic cause of your ICC. The results of a genetic screen may not change anything for you in terms of your diagnosis and management - these decisions are usually based on the results of clinical tests and examination.

However, if the causative spelling mistake is found, it can be a very useful and efficient tool for finding out who else in your family may be at risk of developing the same ICC as you.



There are several possible results from 'genetic screening':

- A genetic 'spelling mistake' is found which is believed to be responsible for causing your ICC. This allows for pre-symptomatic genetic testing of family members to be offered.
- 2) The responsible genetic 'spelling mistake' is not found. This does not mean you do not have the ICC you were clinically diagnosed with, or that it is not an inherited condition, just that the specific responsible genetic 'spelling mistake' in you has not been identified.

This may be because the testing procedure was unable to identify your particular gene alteration or because the particular gene alteration is in genes that the laboratory does not test, or have not been discovered yet. In this situation, pre-symptomatic genetic testing cannot be offered to other family members. However, usual practice is to store your sample, in case additional testing becomes possible in the future.

3) A genetic 'spelling mistake' is found, but it is not yet certain whether this is what is causing your ICC. In this situation, further genetic tests and clinical investigations of your family may be needed to learn more about this gene change.

Pre-symptomatic genetic testing

If an ICC-causing gene change is found in a family member who has already been diagnosed with an ICC (see genetic screening section), pre-symptomatic genetic testing can be offered to blood relatives who are not already known to have the ICC. In the first instance this would usually mean offering testing to close relatives such as parents, brothers, sisters and children, depending on their age.

If a relative chooses to have pre-symptomatic testing, there are two possible outcomes:

- The causative gene change is found. That person is at increased risk of developing the ICC and should have their heart checked by a cardiologist. If they have children, they too would be at risk of the familial ICC and genetic testing and/or screening would be recommended, depending on their age.
- 2) The causative gene change is not found. That person is not at increased risk of developing the ICC and does not need heart checks. If they have children, then they do not need genetic testing or heart checks either.

This approach of finding out who else in the family may have inherited the ICC is sometimes called 'cascade screening'.

The aim of pre-symptomatic testing is to predict someone's future risk of developing the ICC that is known to run in the family. By identifying family members who also have the genetic 'spelling mistake', it is hoped that their risk of ill-health and dangerous heart problems can be reduced through screening, surveillance and appropriate medical and/or surgical therapy.

If you would like to find out more about genetic testing, you should ask your GP or cardiologist to refer you to a specialist inherited cardiac conditions clinic or your local clinical genetics department. You should be offered a genetic counselling appointment with a specialist health professional (usually a genetic counselor or doctor) to discuss genetic testing in greater depth.

If, for any reason, you do not want to have presymptomatic genetic testing you can still have clinical tests such as EKG and echocardiography. Your GP should refer you to a specialist inherited cardiac conditions clinic for this.



What is genetic counselling?

Your genetic counselling appointment is a chance to learn more about the genetic basis of your ICC and to discuss what having the condition may mean for you and your family. You will be able to discuss the pros and cons of genetic testing in reference to your own personal situation, so you can work out whether having a genetic test really is the right choice for you.

Genetic counselling has an especially important role when you are considering pre-symptomatic genetic testing. Discovering you are at increased risk of an ICC when you feel well can have a significant impact on your life, and people can often react to this in different ways. It is worth thinking about your motivation for having genetic testing and finding out what genetic testing can actually tell you. For example, pre-symptomatic genetic testing will only be able to tell you whether you are at an increased risk of developing the ICC. It will not be able to tell you if you will ever develop symptoms, when you will develop them or how severe they may be.

If your pre-symptomatic genetic test shows that you do carry the 'spelling mistake', an outpatient appointment with a heart specialist (cardiologist or specialist nurse) is recommended and this can have implications for insurance, exercise, lifestyle and career choice. So it is worth thinking about how it may affect you and your lifestyle, before and during your appointment.

A genetic counselling appointment will typically last 30-45 minutes and can be used to discuss a variety of issues, as mentioned above, as well as any other questions you may have. The genetic counselor or doctor will draw a family tree and discuss any health issues in the family. You should therefore come prepared with as much relevant personal and family information as possible (including documents where available).

How is the genetic test done?

The genetic test itself involves you giving a blood sample or cheek swab. This will then be sent away to a specialist genetics laboratory.

If it is a genetic screening test, scientists there will look at the genes in your sample to see if they can find a gene change that might be responsible for causing your ICC. They will only look in the genes they know are involved in causing your ICC, not in any other genes, so it will not usually be possible to tell whether you are also at increased risk of other inherited conditions. The results of genetic screening can take several months (sometimes longer) to come through as this is a complex process.

If it is a pre-symptomatic genetic test, the scientists in the lab simply look to see whether you carry the same gene change as your relative(s) or not. The results for pre-symptomatic genetic tests therefore take less time to come back – usually about a month.

Results will be sent back to the genetic counselor or doctor you saw originally and they will then let you know by whatever means you arranged e.g. another appointment, telephone, letter or email. When your genetic counselor or doctor informs you of your results, you can have another discussion as to what are the appropriate next steps to take and ask any more questions that you may have.

Where can I get more information?

If you would like more information about testing then please ask your GP to refer you to either a specialist inherited cardiac conditions clinic, a specialist cardiologist or your local clinical genetics department.

Your genetics consultant/counselor will be able to give more extensive advice. Long QT Syndrome is a rare condition. Experts would suggest that approximately 1 in 2000 people are affected but, as it is often never diagnosed, this is only an estimate. Long QT is a syndrome which can cause a disturbance in the electrical system of the heart, while the mechanical function of the heart can remain completely normal. It is often the result of inheriting an abnormal gene which causes an imbalance in molecules that control the electrical impulses of the heart. Long QT may result in a very fast abnormal heart rhythm (known as an arrhythmia). This particular arrhythmia is technically known as 'Torsade de Pointes'. When this abnormal rhythm occurs the heart loses its normal action and the blood is pumped out. The brain quickly becomes deprived of oxygen, resulting in a loss of consciousness (syncope) and, very occasionally death.

Arrhythmia in patients with LQTS may be triggered by exercise or stressful situations. Not everyone who has LQTS will have an arrhythmia, but if it does occur it can potentially be fatal. LQTS is an electrical problem so it does not change the shape and function of the heart muscle.

Symptoms

It is estimated that up to 50% of people with LQTS do not have symptoms. They may be aware of their condition only from results of an EKG performed for an unrelated reason, because they have a family history of LQTS or because of genetic testing results.

People with LQTS may start to experience symptoms in childhood, although this is not always the case. Symptoms can include unexplained seizures with no warning.

Sudden, unexplained fainting, particularly when in response to a stressful situation. This can often be misdiagnosed as having a 'hysterical reaction'.

Unexplained seizures. A sudden loss of consciousness may be mistaken or misdiagnosed as an epileptic seizure. Sudden cardiac arrest or death in the absence of any structural heart disease or other cardiac problems. Approximately 1 in 10 sudden cardiac arrests or death are the first sign of LQTS.

Swimming in cold water and diving can be particularly dangerous for a person with Long QT, swimming should never be undertaken unaccompanied.

Another sign or symptom of LQTS, is sudden loss of consciousness following a startle or a sudden loud noise.

Causes

LQTS can be inherited or acquired. Acquired LQTS is usually due to the administration of certain medications. Many groups of common medications can cause a prolonged QT interval including certain antibiotics, antihistamines, antidepressants, antipsychotics and heart medications. A comprehensive list of medications can be obtained from your physician or on Google "Long QT drugs". Inherited LQTS is caused by mutations of certain genes and can be passed onto family members. The frequency of inherited LQTS is not known. There are several different types of inherited LQTS and your cardiologist may be able to tell you which type you have. The three most common types of inherited LQTS are called LQTS 1, LQTS 2 and LQTS 3.

At least 17 genes associated with LQTS have been discovered so far, and hundreds of mutations within these genes have been identified. Mutations in three of these genes (LQTS 1, 2 and 3) account for about 70 to 75 percent of LQTS cases. The type of LQTS may be identified by genetic testing. In types 1 and 2, the potassium channels within the heart cause the problem. In these types, arrhythmia may be triggered by exercise or by emotional stress. In type 3 it is the sodium channel that is affected and a low heart rate during sleep or rest may be the trigger for arrhythmia.

Diagnosis

EKG

An EKG is a simple tracing of the heart's electrical activity. It involves attaching electrodes to the chest and limbs and a recording is made. It may reveal a long QT interval which may suggest that it is more likely that you have LQTS. Not all people with LQTS have a prolonged QT interval on their resting EKG and it may be necessary to undertake several EKGs over a period of time or have a period of continuous monitoring using a portable heart monitor.

Exercise tolerance test

Some people may only have a prolonged QT interval when exercising so it may be necessary to have EKG monitoring done while exercising on a treadmill. This is normally the best way for your doctor to find out if you have LQTS. It can also help your doctor find out which type of LQTS you might have.

Neurological test

An electroencephalogram (EEG) test looks for neurological causes of fainting, such as a seizure disorder. The procedure measures the waves of electrical activity the brain produces. Small electrodes attached to your head pick up the electrical impulses from your brain and send them to the EEG machine, which records brain waves. An EEG looks for other conditions that LQTS may be misdiagnosed as, e.g. epilepsy.

Treatment

The main aim in treatment is to prevent loss of consciousness and a life threatening arrhythmia from occurring. There is no cure for LQTS but treatment options include: medications, medical devices, surgery or lifestyle changes. Treatment will be dictated by what type of long QT you have and what is most suitable for you.

It is wise to inform other people if you have LQTS so that they know to call for urgent medical help if you were to faint. Identity bracelets are available from certain charities which carry medical information about you. Your local arrhythmia nurse or your cardiologist may be able to give you more information about this. Alternatively, ask Arrhythmia Alliance for further details.

There are many medications which might affect the heart rhythm in patients with LQTS. These include some over-the-counter cough or cold remedies (decongestants) and some antibiotics. Other drugs that might affect the QT interval include some antidepressants, some treatments for fungal infections, and drugs for heart rhythm disorders.

If you are prescribed any medicines, always check with your doctor and pharmacist that it is safe for a patient with LQTS to take these medicines. Some herbal remedies are also to be avoided (e.g. St John's Wort). Please take care with herbal remedies and ask your doctor for specific advice.



A list of drugs currently known to affect long QT are available now on www.crediblemeds.org. This list will not be exhaustive as newer drugs are becoming more available. Always inform anyone who is prescribing you medication that you have LQTS as there may be newer drugs on the market which may have not have been added to the website.



Medications

Beta blockers: These drugs slow the heart rate and make the dangerous rhythm associated with LQTS less likely. They work by blunting the way a LQTSaffected heart reacts to adrenaline in times of stress, fear or exertion.

Mexiletine: In people with a form of LQTS called LQT₃, taking this antiarrhythmic drug in combination with a beta blocker may help shorten the QT interval.

Potassium: Potassium is a mineral in your body that is important for your heart's electrical system. Potassium supplements may improve the heart's recharging system and may be helpful for people with certain forms of LQTS.

Medical devices and surgery

A pacemaker or implantable cardioverter defibrillator (ICD): These devices are implanted under the skin of your chest and stop a potentially fatal arrhythmia. A pacemaker produces electrical impulses to treat an abnormal heart rhythm. An ICD continuously monitors your heartbeat and will deliver electrical shocks to restore a normal heart rhythm when necessary.

Left cardiac sympathetic denervation surgery: Specific nerves in your chest controlling your heart rhythm are surgically removed which significantly reduces the risk of sudden death. This surgery is generally reserved for people considered at high risk of sudden death, people who do not tolerate the medications or have a fainting spell despite their medications.

WARNING

Recreational drugs such as ecstasy and cocaine are particularly dangerous in patients with Long QT syndrome and CAN BE FATAL. Patients with even mild Long QT syndrome should NEVER experiment with these drugs.

Brugada syndrome is a rare inherited heart rhythm disorder in which the heart is structurally normal, but patients may be at risk of developing a fast heart rhythm due to changes within the ion channels of the heart. Brugada syndrome restricts the flow of sodium ions into the cells of the heart. These ion channels alter the chemical balance of cardiac cells, by adjusting the amount of electrical charge to them. Therefore, if the electrical properties of a cell are faulty this can result in a disturbance of the heart rhythm (arrhythmia).

Symptoms

Signs and symptoms that could mean you have Brugada syndrome include fainting (syncope), irregular heartbeats, fast and chaotic heartbeats and rarely, sudden cardiac arrest.

Diagnosis

Some genes for Brugada syndrome have been identified but the list is not complete. It is therefore impossible to be sure that a patient does not have Brugada syndrome even if a genetic screening, with a blood test or mouth swab, is negative.

If your doctor suspects that you may have Brugada syndrome he will advise you to have a simple test known as the flecainide (or Ajmaline test) challenge to confirm your diagnosis. Flecainide is a drug which blocks sodium channels. As it blocks the faulty sodium channels it unmasks EKG changes in patients with Brugada syndrome. Your doctor will administer the drug through a vein in your hand and record your EKG. The EKG will record how your heart reacts to the flecainide allowing the doctor to collect detailed information about the cause of your potential arrhythmia.

Treatments

If the test result is negative, your doctor will consider your individual risk, and advise you if further tests are necessary. It is likely that you will be able to go home the same day. However, it is important to ask a friend or family member to collect you and drive you home. It is also recommended that you have someone with you for the rest of the day after the test.

If the test is positive, and you are at risk of a fast heart rhythm developing, your doctor may suggest an electrophysiology study and ultimately you may be advised to have an implantable cardioverter defibrillator (ICD) fitted. An ICD will not prevent the arrhythmia but can treat it when one happens. If the test result is positive, it is likely that you will be advised to remain in hospital until after these further tests.



Following your discharge from the hospital you will be able to resume your normal daily activities, including returning to work. It is important to avoid binge drinking alcohol.

Testing for family and relatives

As noted above, genetic testing is at a very early stage for the diagnosis of inherited heart rhythm disorders. However, you and your family may be offered genetic testing if Brugada syndrome is diagnosed in a relative. If the faulty gene is discovered in the suspected case, and then also found in the relative, there is a risk that the relative could suffer symptoms because of Brugada syndrome. If the genetic test is negative, it may not rule out a genetic heart rhythm disorder.

Catecholaminergic Polymorphic Ventricular Tachycardia (CPVT)

Catecholaminergic Polymorphic Ventricular Tachycardia (CPVT) is a rare inherited arrhythmia (heart rhythm disorder). CPVT is sometimes known as a channelopathy. This is because CPVT is caused by an abnormality of ion channels which allow calcium to enter the heart muscle cell. This can lead to an abnormally fast and irregular arrhythmia, ventricular tachycardia – usually during exercise or when adrenaline is released in stressful situations.

These extra beats most frequently arise from the bottom pumping chambers of the heart (ventricular ectopic beats). If there are runs of fast beats from the atria or ventricles these are known as atrial ventricular tachycardia.

In CPVT the heart usually has a normal structure. It is thought that about 1 in 10,000 people have CPVT, although the exact numbers are unknown.

Symptoms of CPVT

Episodes of ventricular tachycardia (VT) can cause light-headedness, dizziness and loss of consciousness (syncope). Symptoms usually start in childhood but can appear in young adults for the first time. Most syncopal episodes in childhood are benign, however those who suffer syncope during exercise or in response to an adrenergic stimulus should be further investigated.



Sometimes these episodes can be mistaken for a fit/epilepsy as they can look very similar and as a consequence many children are treated with antiepileptic drugs. A child not responding to antiepileptic drugs should be referred to a cardiologist for further assessment. Unfortunately, sometimes the first presentation can be sudden cardiac arrest – an episode of VT cannot be sustained for a long period of time and may result in the heart completely stopping (cardiac arrest).



Diagnosis of CPVT

In patients presenting with sudden cardiac arrest in the absence of structural cardiac disease, CPVT should be considered in the differential diagnosis. Clinical diagnosis is made based on family history, exercise or emotional stress induced symptoms and, significantly, response to exercise or catecholamine (adrenaline) infusion, however not all episodes are triggered by adrenaline. The cardiac ultrasound test (echocardiogram) and resting EKG are usually normal.

Genetics

CPVT can result from mutations in the cardiac ryanodine receptor gene in about 50% of cases. Other rarer causes are faulty genes in Calsequestrin, calmodulin and Triadin. Normally, heart muscle cells (myocytes) contract and relax in a coordinated way, but mutations in either RYR2 or CASQ2 impairs calcium handling within these myocytes and so during exercise or emotional stress, ventricular tachycardia may result. Familial occurrence has been seen in about a third of cases of CPVT. RYR2 causes autosomal dominant inheritance meaning if you inherit the abnormal gene from only one parent, you can get the disease. CASQ2 is inherited in an autosomal recessive manner meaning two copies of an abnormal gene must be present in order for CPVT to develop.

Currently, genetic testing is not widely available. However first-degree relatives should be evaluated with ECG, Holter monitoring and exercise stress testing. Genetic testing can sometimes be helpful. Identification of a genetic abnormality that is the definite cause of the condition (a "pathogenic" mutation) can allow other family members to be tested. Genetic analysis might identify silent carriers of CPVT– related mutations, and it may be recommended that even symptom-free carriers are treated with medication such as beta blockers. Sometimes, a genetic change is discovered, and it is unclear if this is the definite cause. This is known as a "variant of unknown significance" or VUS. A VUS cannot be used for family screening.



Treatment

Once diagnosed, treatment is usually with a beta blocker. Beta blockers decrease the activity of the heart by blocking the action of hormones such as adrenaline, which would normally increase in times of exercise or emotional stress. Subsequently, the number of episodes of VT is reduced. A high dose is often required. Flecainide may also be used in addition to beta blockers if the response is inadequate. Flecainide inhibits cardiac ryanodine receptor-mediated calcium release.

Missing even a single dose of beta blocker can be potentially dangerous. Internal cardiac defibrillators (ICD) are sometimes fitted in addition to medication to 'shock' the heart back into normal rhythm if VT occurs. Survivors of cardiac arrest or high-risk patients with a strong family history of sudden death are likely to receive an ICD. ICD treatment without use of beta blockers is not advised as a shock from the defibrillator can lead to an adrenaline surge and multiple runs of ventricular tachycardia known as an 'electrical storm'.

A left cervical sympathectomy may be offered to some patients e.g. those in whom beta blockers are contraindicated, when an ICD cannot be fitted, or where there is recurrent VT in patients with an ICD despite maximal medical treatment. A cervical sympathectomy is an operation carried out through a small incision under the arm. This blocks a group of nerves that produce and deliver adrenaline to the heart. These nerves are not essential to normal heart function but sympathectomy can be very helpful in preventing serious arrhythmias.

Additionally, some sports may be restricted, e.g. swimming, and certain medications that increase the heart rate may need to be avoided.

Notes

| |
|------|
| |
| |

This booklet has been written to support those diagnosed with an arrhythmia and their carers, who struggle to find information on this condition. Without donations and fundraising, we would not be able to provide support through our award-winning resources and helpline.



Working together to improve the diagnosis, treatment and quality of life for all those affected by arrhythmias



Arrhythmia Alliance

843-415-1886

info-us@heartrhythmalliance.org
www.heartrhythmalliance.org

Registered Non-Profit 501(c)(3) ©Arrhythmia Alliance

Published 2005 Reviewed April 2023



Finger on your Pulse: is our new library of educational video resources. Medical Experts share their knowledge and address specific concerns and patients share their experience living with the various conditions and treatments.

www.fingeronyourpulse.org

"it is important to be aware of a genetic risk within your family, The booklet advised me to ensure both my daughter's were checked for LQTS, It was such a relief to know they were in the clear"

Carl

Please remember that this publication provides general information. You should always discuss and seek advice from your healthcare professional as to what is most appropriate for you.

Acknowledgments: Arrhythmia Alliance would like to thank all those who helped in the development and review of this publication. Particular thanks are given to Dr Charlotte D'Souza and Dr Jan Till.

Founder and Trustee:

Trudie Lobban MBE, FRCP (Edin)